

Patient Identification	Prescriber Details
Name:	Name:
	Signature:
	Date:

## Prescriber's Checklist for Women of Childbearing Potential

Prior to Initiation of Treatment
<p>Treatment with REBLOZYL should not be started if the woman is pregnant or in women of childbearing potential not using an effective method of contraception.</p> <ul style="list-style-type: none"> <li>• The use of REBLOZYL is not recommended in pregnancy and in women of childbearing potential not using effective contraception.</li> <li>• There are no data from the use of REBLOZYL in pregnant women. Studies in animals have shown reproductive toxicity and embryo-foetal toxicity. Clinical implications are potential foetal loss and teratogenicity.</li> </ul>
<input type="checkbox"/> Provide counselling before treatment initiation regarding the potential teratogenic risk of REBLOZYL and required actions that may help to minimize this risk.
<input type="checkbox"/> Inform women of childbearing potential of the necessity for an effective method of contraception while on treatment and for 3 months after discontinuation.
<input type="checkbox"/> A pregnancy test is recommended in women of childbearing potential before starting treatment.
<input type="checkbox"/> Provide the Patient Card to women of childbearing potential.

Duration of Treatment
<input type="checkbox"/> Provide regular counselling regarding the potential teratogenic risk of REBLOZYL and required actions that may help to minimize this risk.
<input type="checkbox"/> Remind women of childbearing potential that they must use an effective method of contraception during treatment with REBLOZYL.
<p>During treatment with REBLOZYL, women must not become pregnant. If a woman becomes pregnant or wants to become pregnant, she should be informed of the potential for hazard to the fetus.</p>

Discontinuation of Treatment
<input type="checkbox"/> Counsel women of childbearing potential that an effective method of contraception should be maintained for at least 3 months following discontinuation of treatment with REBLOZYL.
<input type="checkbox"/> Provide counselling in the event of pregnancy and evaluation of the outcome of any pregnancy.
<input type="checkbox"/> Not applicable (this patient did not become pregnant while on treatment or within 3 months of discontinuation of REBLOZYL)

**Should a pregnancy occur during treatment or within 3 months following discontinuation of treatment with REBLOZYL, remind the patient that it should be reported to the prescriber.**

Celgene Inc. encourages healthcare professionals and patients to report any pregnancy and any suspected adverse reactions. This will allow quick identification of new safety information. More information regarding how to report side effects can be found in the Canadian Product Monograph: [https://www.bms.com/assets/bms/ca/documents/productmonograph/REBLOZYL\\_EN\\_PM.pdf](https://www.bms.com/assets/bms/ca/documents/productmonograph/REBLOZYL_EN_PM.pdf). For more information or to obtain a copy of this document, please contact medical information by calling 1-866-463-6267.

**CLEAR FORM**